




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 309/38, 309/40, 405/04, 409/04,</b> <b>A61K 31/35, C07C 317/24, 311/29</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/18753</b> <b>(43) International Publication Date:</b> 6 April 2000 (06.04.00)
<b>(21) International Application Number:</b> PCT/EP99/06873 <b>(22) International Filing Date:</b> 16 September 1999 (16.09.99) <b>(30) Priority Data:</b> P 9802011 25 September 1998 (25.09.98) ES P 9900619 26 March 1999 (26.03.99) ES <b>(71) Applicant (for all designated States except US):</b> ALMIRALL PRODESFARMA, S.A. [ES/ES]; General Mitre 151, E-08022 Barcelona (ES). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> CRESPO CRESPO, M <sup>a</sup> Isabel [ES/ES]; Comte d'Urgell 259, 6 <sup>o</sup> -3 <sup>a</sup> , E-08036 Barcelona (ES). JIMENEZ MAYORGA, Juan Miguel [ES/ES]; Dr. Roux 74, Ppal.2 <sup>a</sup> , E-08017 Barcelona (ES). PUIG DURAN, Carles [ES/ES]; Asturias, 99, 2 <sup>o</sup> -2 <sup>a</sup> , E-08024 Barcelona (ES). SOCA PUEYO, Lidia [ES/ES]; Burdeos 2, 3 <sup>o</sup> 1 <sup>a</sup> , E-08029 Barcelona (ES). <b>(74) Agent:</b> GOLDIN, Douglas, Michael; J.A. Kemp & Co. 14 South Square, Gray's Inn, London WC1R 5LX (GB).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>

**(54) Title:** 2-PHENYLPYRAN-4-ONE DERIVATIVES

(I)

**(57) Abstract**

2-Phenylpyran-4-one derivatives of formula (I), wherein R<sup>1</sup> represents an alkyl or -NR<sup>4</sup>R<sup>5</sup> group, wherein R<sup>4</sup> and R<sup>5</sup> each independently represents a hydrogen atom or an alkyl group; R<sup>2</sup> represents an alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, pyridyl, thienyl, naphthyl, tetrahydronaphthyl or indanyl group, or a phenyl group which may be unsubstituted or substituted by one or more halogen atoms or alkyl, trifluoromethyl, hydroxy, alkoxy, methylthio, amino, mono- or dialkylamino, hydroxyalkyl or hydroxycarbonyl groups; R<sup>3</sup> represents a methyl, hydroxymethyl, alkoxymethyl, C<sub>3</sub>-C<sub>7</sub> cycloalkoxymethyl, benzyloxymethyl, hydroxycarbonyl, nitrile, trifluoromethyl or difluoromethyl group or a CH<sub>2</sub>-R<sup>6</sup> group wherein R<sup>6</sup> represents an alkyl group; and X represents a single bond, an oxygen atom, a sulfur atom or a methylene group; or pharmaceutically acceptable salts thereof, processes for their production and synthetic intermediates used in said processes, pharmaceutical compositions containing them and their use in medical treatment.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		


## 2-PHENYLPYRAN-4-ONE DERIVATIVES

This invention relates to new therapeutically useful 2-phenylpyran-4-one derivatives, to processes for their preparation and to pharmaceutical compositions containing them.

It is known that non-selective inhibition of the enzyme cyclooxygenase (COX) prevents the overproduction of prostaglandins associated with inflammation, which are mediated by cyclooxygenase-2 (COX-2) but, at the same time, deprives tissues of basal levels of prostaglandins necessary for the health of certain tissues mediated largely by cyclooxygenase-1 (COX-1). Non steroidal anti-inflammatory drugs are non-selective inhibitors of COX and for that reason, have side effects of decreased renal blood flow, decreased platelet function, dyspepsia and gastric ulceration.

We have now found that certain 2-phenylpyran-4-one derivatives selectively inhibit COX-2 in preference to COX-1 and are useful in the treatment of COX-2 mediated diseases, such as inflammation, pain, fever and asthma with fewer side effects.

Accordingly the present invention provides a 2-phenylpyran-4-one compound of formula (I):



(I)

wherein:

$\text{R}^1$  represents an alkyl or  $-\text{NR}^4\text{R}^5$  group, wherein  $\text{R}^4$  and  $\text{R}^5$  each independently represents a hydrogen atom or an alkyl group;

$\text{R}^2$  represents an alkyl,  $\text{C}_3\text{-C}$ , cycloalkyl, pyridyl, thienyl, naphthyl, tetrahydronaphthyl or indanyl group, or a phenyl

group which may be unsubstituted or substituted by one or more halogen atoms or alkyl, trifluoromethyl, hydroxy, alkoxy, methylthio, amino, mono- or dialkylamino, hydroxyalkyl or hydroxycarbonyl groups;

5         $R^3$  represents a methyl, hydroxymethyl, alkoxymethyl,  $C_3$ - $C_7$  cycloalkoxymethyl, benzyloxymethyl, hydroxycarbonyl, nitrile, trifluoromethyl or difluoromethyl group or a  $CH_2$ - $R^6$  group wherein  $R^6$  represents an alkyl group; and

10        X represents a single bond, an oxygen atom, a sulfur atom or a methylene group;

or a pharmaceutically acceptable salt thereof.

The alkyl groups and moieties such as those present in the alkoxy, hydroxyalkyl, mono- or di-alkylamino groups, mentioned in relation to the groups  $R^1$  to  $R^6$  are usually "lower" alkyl  
15 that is containing from 1 to 6 particularly from 1 to 4 carbon atoms, the hydrocarbon chain being branched or straight. Preferred alkyl groups, and where relevant alkyl moieties, include methyl, ethyl, propyl including i-propyl, and butyl including n-butyl, t-butyl and sec-butyl.

20        In a phenyl group substituted by one or more halogen atoms or alkyl, trifluoroalkyl, hydroxy, alkoxy, methylthio, amino, mono- or dialkyl amino, hydroxyalkyl or hydroxycarbonyl groups, the phenyl ring may be substituted by 1, 2, 3, 4 or 5 substituents, preferably 1, 2 or 3 substituents, each being  
25 independently selected from the possible substituents set out above. The phenyl group (attached to X or the pyran-4-one ring through its 1-position) may be substituted at any of the remaining positions, that is to say the 2, 3, 4, 5 or 6-positions. A phenyl group having more than one substituent may  
30 be substituted at any combination of positions. For example a phenyl group having two substituents may be substituted at the 2 and 3, 2 and 4, 2 and 5, 2 and 6, 3 and 4 or 3 and 5 positions.

In particular, it is preferred that  $R^2$  represents a  
35 branched alkyl,  $C_3$ - $C_7$  (preferably  $C_3$ ,  $C_5$  or  $C_6$ ) cycloalkyl, naphthyl, tetrahydronaphthyl or indanyl group, an unsubstituted phenyl group or a phenyl group substituted by one or more

halogen atoms, alkoxy groups, preferably methoxy groups, and/or alkyl groups, preferably methyl groups. The phenyl group preferably has 1, 2 or 3 substituents, more preferably 1 or 2 substituents. Halogen atoms are preferably selected from  
5 fluorine, chlorine and bromine atoms. When  $R^2$  is a phenyl group substituted by one or more halogen atoms, alkoxy groups and/or alkyl groups, preferably one of the substitutions is at the 4-position of the phenyl group. When  $R^2$  is a phenyl group substituted by one or two halogen atoms at least one of the  
10 substitutions is preferably on the 2- or the 4-position.

It is preferred that  $R^1$  independently represents an unsubstituted alkyl group such as methyl, ethyl, propyl or butyl, preferably methyl, or an  $NH_2$  group (i.e.  $R^4$  and  $R^5$  in the above formula both independently represent an H atom).

15 It is also preferred that  $R^3$  independently represents an unsubstituted alkyl group such as methyl, ethyl, propyl or butyl, preferably methyl, a nitrile group, a hydroxymethyl group, a methoxymethyl group, a difluoromethyl group or a hydroxycarbonyl group.

20 It is further preferred that X independently represents a single bond, an oxygen atom or a methylene group more preferably a single bond or an oxygen atom.

Specific examples of the 2-phenylpyran-4-one derivatives of the present invention include:

25 2-(4-methanesulfonylphenyl)-6-methyl-3-phenylpyran-4-one,  
3-(4-fluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(3-fluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
30 3-(2-fluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(4-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(3-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-  
35 one,  
3-(2-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,

- 3-(4-bromophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
2-(4-methanesulfonylphenyl)-6-methyl-3-*p*-tolylpyran-4-one,  
2-(4-methanesulfonylphenyl)-6-methyl-3-*m*-tolylpyran-4-one,  
5 2-(4-methanesulfonylphenyl)-6-methyl-3-*o*-tolylpyran-4-one,  
2-(4-methanesulfonylphenyl)-6-methyl-3-(4-trifluoromethylphenyl)pyran-4-one,  
3-(2,4-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
10 3-(3,4-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(3,5-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(2,5-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
15 3-(2,6-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(2,4-dichlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
20 3-(3,4-dichlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(3-fluoro-4-methoxyphenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(4-chloro-3-fluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
25 3-(2-chloro-4-fluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(4-bromophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
30 3-(4-fluorophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(2,4-difluorophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-cyclohexyl-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
35 2-(4-methanesulfonylphenyl)-6-methyl-3-naphthalen-2-ylpyran-4-one,


- 4- (6-methyl-4-oxo-3-phenyl-4H-pyran-2-yl) benzenesulfonamide,  
4- [3- (4-fluorophenyl) -6-methyl-4-oxo-4H-pyran-2-yl]  
benzenesulfonamide,  
4- [3- (3,4-dichlorophenyl) -6-methyl-4-oxo-4H-pyran-2-yl]  
5 benzenesulfonamide,  
5- (2,4-difluorophenyl) -6- (4-methanesulfonylphenyl) -4-oxo-4H-  
pyran-2-carbonitrile.  
3- (2-fluorophenoxy) -2- (methanesulfonylphenyl) -6-methylpyran-4-  
one,  
10 3- (4-chlorophenoxy) -2- (methanesulfonylphenyl) -6-methylpyran-4-  
one,  
3- (2-chlorophenoxy) -2- (methanesulfonylphenyl) -6-methylpyran-4-  
one,  
3- (2,5-difluorophenoxy) -2- (methanesulfonylphenyl) -6-  
15 methylpyran-4-one,  
3- (3-chloro-4-methylphenyl) -2- (4-methanesulfonylphenyl) -6-  
methylpyran-4-one,  
2- (4-methanesulfonylphenyl) -6-methyl-3-phenoxy pyran-4-one,  
3- (4-fluorophenoxy) -2- (4-methanesulfonylphenyl) -6-methylpyran-  
20 4-one,  
2- (4-methanesulfonylphenyl) -6-methyl-3- (4-methylphenoxy) pyran-  
4-one,  
3- (4-chlorophenyl) -2- (4-methanesulfonylphenyl) -6-methoxy-  
methylpyran-4-one,  
25 3- (4-chlorophenyl) -6-difluoromethyl-2- (4-methanesulfonyl-  
phenyl) pyran-4-one,  
and anyone of the compounds specifically identified in Table 4,  
and pharmaceutically acceptable salts thereof.
- 30 Of outstanding interest are:  
3- (4-fluorophenyl) -2- (4-methanesulfonylphenyl) -6-methylpyran-4-  
one,  
3- (2-fluorophenyl) -2- (4-methanesulfonylphenyl) -6-methylpyran-4-  
one,  
35 3- (4-chlorophenyl) -2- (4-methanesulfonylphenyl) -6-methylpyran-4-  
one,

- 3-(4-bromophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(2,4-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
5 3-(3,4-dichlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(3-chloro-4-methylphenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
2-(4-methanesulfonylphenyl)-6-methyl-3-phenoxy-pyran-4-one,  
10 3-(4-fluorophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(2-fluorophenoxy)-2-(methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(4-chlorophenoxy)-2-(methanesulfonylphenyl)-6-methylpyran-4-one,  
15 one,  
3-(2-chlorophenoxy)-2-(methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(4-bromophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
20 2-(4-methanesulfonylphenyl)-6-methyl-3-(4-methylphenoxy) pyran-4-one,  
3-(2,4-difluorophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(2,5-difluorophenoxy)-2-(methanesulfonylphenyl)-6-methylpyran-4-one,  
25 3-(4-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methoxymethylpyran-4-one,  
3-(4-chlorophenyl)-6-difluoromethyl-2-(4-methanesulfonylphenyl)pyran-4-one,  
30 and pharmaceutically acceptable salts thereof.

The present invention also provides processes for preparing a compound of formula (I) which depend on the definition of  $R^3$ . When  $R^3$  is a methyl group, compounds of formula (I) are prepared according to the definition of  $R^1$ .


- 35 Thus, compounds of formula (I) in which  $R^3$  is a methyl group and  $R^1$  is an alkyl or  $-NR^4R^5$  group in which  $R^4$  and  $R^5$  are alkyl groups, viz. a 2-phenylpyran-4-one derivative of formula (II):





(II)


10 wherein  $R^{1a}$  is an alkyl or  $-NR^{4a}R^{5a}$  group in which  $R^{4a}$  and  $R^{5a}$  are each independently alkyl groups, and  $R^2$  and  $X$  are as defined above, which comprises reacting a carbonyl derivative of formula (III):



(III)


15 wherein  $R^{1a}$ ,  $R^2$  and  $X$  are as defined above with an excess of anhydrous acetic acid and polyphosphoric acid at a temperature from 100°C to 150°C.

The carbonyl derivative of formula (III) may be obtained by methods well known in the literature (EP-A-714883; WO96/06840; WO96/31509 and DE-2064520) or when  $X$  represents an oxygen or sulfur atom, by reacting a phenacyl derivative of  
20 formula (IV):



(IV)

wherein  $R^{1a}$  is as defined above and  $Y$  represents a chlorine or bromine atom, with a hydroxy or mercapto derivative of formula  
30 (V):




(V)

wherein  $R^2$  is as defined above and  $X^a$  is an oxygen or sulfur atom.


The reaction between the phenacyl derivative of formula (IV) and the intermediate compound of formula (V) may be carried out by heating a mixture of these two starting materials, optionally in a solvent mixture of methylene chloride, benzene or toluene and water, at a temperature of from 15°C to 30°C and in the presence of a phase transfer catalyst as benzyltriethylammonium chloride.

The carbonyl derivative of formula (III) in which X is other than a sulfur atom, may also be prepared by reacting a thio derivative of formula (VI):




wherein  $R^{1a}$  and  $R^2$  are as defined above, and  $X^b$  is a single bond, an oxygen atom or a methylene group, with an oxidizing agent, preferably magnesium monoperoxyphthalate or 3-chloroperoxybenzoic acid. The reaction is preferably carried out in an organic solvent such as a mixture of methylene chloride with methanol or ethanol, at a temperature of from 10°C to 40°C.

The present invention also provides a process for the preparation of a compound of formula (I) wherein  $R^3$  is a methyl group,  $R^1$  is an alkyl group, and X is other than a sulfur atom viz. 2-phenylpyran-4-one derivative of formula (VII):




wherein  $R^{1b}$  is an alkyl group and  $R^2$  and  $X^b$  are as defined above by reacting a mercapto derivative of formula (VIII):




10 wherein  $R^{1b}$ ,  $R^2$  and  $X^b$  are as defined above with an oxidizing agent, preferably with magnesium monoperoxyphthalate or 3-chloroperoxybenzoic acid.

The reaction between the mercapto derivative of formula (VIII) and the oxidizing agent is preferably carried out, as previously disclosed for the compound of formula (VI), in an organic solvent such as a mixture of methylene chloride with methanol or ethanol, at a temperature of from 10°C to 40°C.

The present invention additionally provides a process for the preparation of a compound of formula (I) wherein  $R^1$  is a  $-NR^4R^5$  group and  $R^3$  is a methyl group, viz. 2-phenylpyran-4-one derivative of formula (IX):



30 wherein  $R^2$ ,  $R^4$ ,  $R^5$  and  $X$  are as defined above by reacting a chlorosulfonyl derivative of formula (X):




wherein  $R^2$  and X are as defined above with an amine of formula (XI):



wherein  $R^4$  and  $R^5$  are as defined above.

This reaction is preferably carried out at a temperature of from 10°C to 40°C.


The chlorosulfonyl derivative of formula (X) may, for example, be prepared by reacting a compound of formula (XII):



(XII)


wherein  $R^2$  and X are as defined above with chlorosulfonic acid, preferably at a temperature of from 80°C to 120°C.

The present invention further provides a process for the preparation of a compound of formula (I) wherein  $R^3$  is a methyl group and  $R^1$  is a  $-NR^4R^5$  group wherein  $R^4$  and  $R^5$  are hydrogen, viz, the 2-phenylpyran-4-one derivative of formula (XIII):



(XIII)

wherein  $R^2$  and X are as defined above by debenzoylation of the corresponding N,N-dibenzoyl derivative of formula (XIV):



(XIV)

wherein R<sup>2</sup> and X are as defined above.

The debenzylation is preferably carried out with an excess of trifluoroacetic, sulfuric or methanesulfonic acid at a temperature of from 0°C to 120°C.


The intermediate of formula (XIV) may be prepared according to the above processes using appropriate starting materials wherein R<sup>i</sup> and R<sup>j</sup> (or R<sup>4a</sup> and R<sup>5a</sup>) both represent benzyl groups.

The intermediate of formula (IV) and (VI) used in the preparation of the compounds of the invention may be prepared by methods disclosed in the literature, for example, in M.F. Saettone, J. Org. Chem. 31, p. 1959 (1966) and WO-9606840.

The intermediate compounds of formulae (VIII) and (XII) may be prepared by the same process disclosed for the preparation of compounds of formula (II), with the appropriate starting materials.

The 2-phenylpyran-4-one derivatives of formula (I) wherein R<sup>3</sup> is other than a methyl group, can be prepared by processes which are represented in the following scheme:

## SCHEME



As can be seen in the scheme, 2-phenylpyran-4-one derivatives of formula (I) wherein  $R^3$  is other than a methyl group, viz. compounds of formulae (XVII), (XVIII), (XIX), (XX), (XXI), (XXII) and (XXIV), are prepared from compounds of formula (I) in which  $R^3$  is a methyl group, viz. compound of formula (XV), which processes of preparation have been disclosed above. In a first stage, compounds of formula (XV) are treated with an oxidizing agent as selenium dioxide in an organic solvent as tetrahydrofuran or dioxan, in a pressure vessel and at a temperature from 100°C to 190°C. The corresponding aldehyde of formula (XVI) is obtained, which is used as starting material to obtain the compounds of formula (I) with  $R^3$  other than a methyl group.

The compounds of formula (I) wherein  $R^3$  is a hydroxycarbonyl group, viz. compound of formula (XVII), are prepared from the corresponding aldehyde (XVI) by reaction with an oxidizing agent as pyridinium dichromate or manganese dioxide in an organic solvent as N,N-dimethylformamide or ethanol at a temperature between -5°C and 35°C. The obtained compounds (XVII) are used as starting materials to obtain compounds of formula (I) wherein  $R^3$  is a trifluoromethyl group, viz. compound of formula (XVIII). The reaction is carried out by reaction of compounds (XVII) with a mixture of sulphur tetrafluoride and hydrogen fluoride, optionally in the presence of an organic solvent as methylene chloride, in a pressure vessel, and at a temperature from 20°C to 140°C.

The compounds of formula (I) wherein  $R^3$  represents a hydroxymethyl group viz. compounds of formula (XIX) are prepared by reduction of compounds (XVI) with a boron or aluminium hydride, preferably sodium borohydride in a solvent as methanol or ethanol and at a temperature from 10°C to 40°C. Further reaction of compounds (XIX) with an appropriate halide of formula (XXIII):

35  $Z - R^7$  (XXIII)

wherein Z represents a chlorine, bromine or iodine atom and  $R^7$

represents an alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl or benzyl group, provide the compounds of formula (I) wherein R<sup>3</sup> is an alkoxymethyl, C<sub>3</sub>-C<sub>7</sub> cycloalkoxymethyl or benzyloxymethyl group viz. compounds of formula (XX). The reaction is carried out in an organic solvent  
5 such as acetone, N,N-dimethylformamide or tetrahydrofuran in the presence of sodium or potassium hydride or amide, and at a temperature between 20°C and 120°C.

Also aldehydes of formula (XVI) are used as starting material to obtain compounds of formula (I) wherein R<sup>3</sup> is a  
10 nitrile group, viz. compounds of formula (XXI). The reaction is carried out in a first stage by treatment of aldehydes (XVI) with hydroxylamine hydrochloride and formic acid at a temperature from 80°C to 120°C. The resulting oxime derivative is isolated and heated with an excess of acetic anhydride at a  
15 temperature between 100°C to 180°C.

The compounds of formula (I) wherein R<sup>3</sup> represents a difluoromethyl group, viz. compounds of formula (XXII), are prepared from aldehydes of formula (XVI) by reaction with a fluorinated reagent as diethylaminosulfur trifluoride or a  
20 mixture of sulfur tetrafluoride-hydrogen fluoride, optionally in the presence of an organic solvent as methylene chloride, benzene or toluene and at a temperature from 0°C to 130°C.

The 2-phenylpyran-4-one derivatives of formula (I) in which R<sup>3</sup> is a CH<sub>2</sub>-R<sup>6</sup> group, viz. compounds of formula (XXIV),  
25 are also prepared from aldehydes of formula (XVI) in a two stages process. In the first stage, the aldehyde (XVI) is reacted with a triphenylphosphine derivative (XXV) in the presence of a solvent as dioxane, dimethoxyethane or tetrahydrofuran at a temperature from 15°C to the boiling point  
30 of the solvent. The resulting compound is hydrogenated in the second stage of the process in the presence of a catalyst as palladium on activated carbon. The reaction is carried out in the presence of a solvent as methanol, ethanol or ethyl acetate at a temperature from 15°C to 40°C.

35 The 2-phenylpyran-4-one derivatives of formula (I) in which there is the presence of a basic group, can be converted by methods known per se into pharmaceutically acceptable salts,



preferably acid addition salts by treatment with organic or inorganic acids as fumaric, tartaric, succinic or hydrochloric acid. Also, 2-phenylpyran-4-one derivatives of formula (I) in which R<sup>3</sup> represents an hydroxycarbonyl group, may be converted  
5 into pharmacologically acceptable salts with, for instance, alkali metals such as sodium or potassium by reaction with an alkali metal hydroxide.

The following biological tests and data further illustrate this invention.

10

#### COX-1 and COX-2 activities in human whole blood

Fresh blood from healthy volunteers who had not taken any non-steroidal anti-inflammatory drugs for at least 7 days prior to blood extraction was collected in heparinized tubes (20  
15 units of heparin per ml). For the COX-1 activity determination, 500  $\mu$ l aliquots of blood were incubated with either 5  $\mu$ l vehicle (dimethylsulphoxide) or 5  $\mu$ l of a test compound for 1 h at 37°C. Calcium ionophore A23187 (25  $\mu$ M) was added 20 min before stopping the incubation. Plasma was separated by  
20 centrifugation (10 min at 13000 rpm) and kept at -30°C until TXB<sub>2</sub> levels were measured using an enzyme immunoassay kit (ELISA). The effect of the compounds were evaluated by incubating each compound at five to six different concentrations with triplicate determinations. IC<sub>50</sub> values were  
25 obtained by non-linear regression using InPlot, GraphPad software on an IBM computer.

For the COX-2 activity determination, 500  $\mu$ l aliquots of blood were incubated in the presence of LPS (10  $\mu$ g/ml) for 24 h at 37°C in order to induce the COX-2 expression (Patriagnani  
30 et al., J. Pharm. Exper. Ther. 271; 1705-1712 (1994)). Plasma was separated by centrifugation (10 min at 13000 rpm) and kept at -30°C until PGE<sub>2</sub> levels were measured using an enzyme immunoassay kit (ELISA). The effects of inhibitors were studied by incubating each compound (5  $\mu$ l aliquots) at five to six  
35 different concentrations with triplicate determinations in the presence of LPS for 24 hours. IC<sub>50</sub> values were obtained by non-linear regression using InPlot, GraphPad software on an IBM

computer.

Anti-inflammatory activity (adjuvant arthritis)

Male Wistar rats weighing 175-200 g with free access to food and water were used. On day 0, the animals received an intraplantar injection of a suspension of Mycobacterium tuberculosis in paraffin oil (0.5 mg/rat) on the left hind paw. A group of nonarthritic control rats were injected with paraffin oil alone. On days 11 and 14 after induction of arthritis, the volume of the hind paw of each rat was measured using a water plethysmograph. Animals whose paw volumes increased during that time were selected. Rats were distributed into groups of 8 having equal mean paw volumes and an approximately equal standard deviation.

Test compounds were administered p.o. once daily for 7 days (days 14-20). Nonarthritic and arthritic control rats received vehicle alone for 7 days. The hind paw volumes were measured 20 h after the last dose (on day 21). The body weight was determined every second day.

The results were expressed as the percentage of inhibition of inflammation (paw volume) for each treatment group, considering both the arthritic and nonarthritic vehicle controls. The ANOVA tests was used for statistical studies.

25 Ulcerogenic activity

Animals : Male Wistar rats (Interfauna, U.K., Ltd.) weighing about 150-175 g were used. Animals were maintained on a 12:12 hour light-dark cycle (lights on at 7:00 am) at room temperature (22±1°C). Food and water were allowed ad libitum.

30

Procedure : The compounds were administered by the oral route once a day for 4 consecutive days. The body weight of each rat was assessed every day before drug administration. The animals were anesthetized 24 h after the last dosing and 1 ml of blood was extracted by cardiac puncture using heparin (10 U/ml) as anticoagulant. The percentage of hematocrit was measured. The intestines were removed, opened longitudinally

and gently washed. The macroscopic severity of the intestinal erosions was assessed using a parametric scale, evaluating the number of the perforated and non-perforated intestinal ulcers by means of a lesion index ranging from 0 to 3 (0: no ulcers, 1:<10 ulcers, 2:10-25 ulcers to 3:>25 ulcers). No gastric ulcers are observed using this protocol.

The treatments were randomized in each experiment. The results were compared with those obtained in the vehicle-treated group using the ANOVA test.

### Results

The results obtained from the biological assays are shown in Table 1, 2 and 3.

TABLE 1: COX-1 and COX-2 Inhibition

Compound (*)	COX-1 IC <sub>50</sub> (μM)	COX-2 IC <sub>50</sub> (μM)	Ratio COX-1/COX-2
Indomethacin	0.19	0.22	0.86
2	>100	1.06	>94
4	>100	1.5	>66
5	>100	2.1	>47
8	>100	1.7	>58
15	100	1.1	90
22	37.1	0.7	53
31	>100	1.67	>59
37	>100	1.08	>92
39	>100	0.96	>104
40	27	0.14	193
41	>100	0.35	>285
42	41	0.2	205
43	>100	0.8	125

Compound (*)	COX-1 IC <sub>50</sub> (μM)	COX-2 IC <sub>50</sub> (μM)	Ratio COX-1/COX-2
44	39	0.21	185
45	22	0.15	147
47	57.1	0.8	71
63	44	1.73	25
67	>100	2.1	>47

(\*) See structures in Table 4.

Indomethacin is 1-(4-chlorobenzoyl)-5-methoxy-2-methylindole-3-acetic acid.

TABLE 2: Anti-inflammatory activity

Compound	% Inhibition (dose, mg/kg)
Indomethacin	64 (1)
5	50 (1)
22	69 (1)
39	75 (1)
41	71 (1)
45	74 (1)

TABLE 3: Ulcerogenic activity

Compound	Dose (mg/kg)	Lesion index		Hematocrit (%)
		PU	NPU	
Vehicle		0	0	44.3±0.2
Indomethacin	10	3	3	22.7±1.6
5	100	0	0	44.1±0.7
22	100	0	0	44.4±0.3

Compound	Dose (mg/kg)	Lesion index		Hematocrit (%)
		PU	NPU	
39	100	0	0	43.7±0.4
41	100	0	0	43.4±1.9
45	100	0	0	44.4±0.9

PU: perforated ulcers, NPU: non-perforated ulcers.

5

As shown in Table 1, the 2-phenylpyran-4-one derivatives of formula (I) are potent and selective COX-2 inhibitors whereas the reference compound indomethacin is as equipotent as COX-2 and COX-1 inhibitor. Due to their low COX-1 inhibitory activity, the compounds of formula (I) present an important anti-inflammatory activity (see Table 2) and the benefit of significantly less harmful side effects than the non-steroidal anti-inflammatory drugs commonly used (e.g. gastrointestinal toxicity (see Table 3), renal side-effects, reduced effect on bleeding times and asthma induction in aspirin-sensitive subjects).

Thus the compounds of the invention are preferably selective inhibitors of mammalian COX-2, for example human COX-2. The compounds of the invention also preferably have low inhibitory activity toward mammalian COX-1, for example human COX-1. Inhibitory activity can typically be measured by *in vitro* assays, for example as described above.

Preferred compounds of the invention have an IC<sub>50</sub> value for COX-2 of less than 5  $\mu$ M, preferably less than 3 more preferably less than 2.5  $\mu$ M. Preferred compounds of the invention also have an IC<sub>50</sub> value for COX-1 of greater than 10  $\mu$ M, preferably greater than 20  $\mu$ M. As an indicator of selectivity for inhibition of COX-2 over COX-1, the ratio of COX-1/COX-2 IC<sub>50</sub> values is preferably greater than 20, 30 or 50, more preferably greater than 80, 90 or 100.

The present invention further provides a compound of formula (I) for use in a method of treatment of the human or animal body by therapy, in particular for the treatment of

pain, fever or inflammation, to inhibit prostanoid-induced smooth muscle contraction or for the prevention of colorectal cancer or neurodegenerative diseases.

The present invention further provides the use of a  
5 compound of formula (I) in the manufacture of a medicament for the treatment of pain, fever or inflammation, to inhibit prostanoid-induced smooth muscle contraction or for the prevention of colorectal cancer or neurodegenerative diseases.

The compounds of formula (I) are useful for relief of  
10 pain, fever and inflammation of a variety of conditions including rheumatic fever, symptoms associated with influenza or other viral infections, common cold, low back and neck pain, dysmenorrhoea, headache, toothache, sprains and strains, myositis, neuralgia, synovitis, bursitis, tendinitis, injuries,  
15 following surgical and dental procedures and arthritis including rheumatoid arthritis, osteoarthritis, gouty arthritis, spondyloarthropathies, systemic lupus erythematosus and juvenile arthritis and bone resorption. They may also be used in the treatment of skin inflammation disorders such as  
20 psoriasis, eczema, burning and dermatitis. In addition, such compounds may be used for the prevention of colorectal cancer.

The compounds of formula (I) will also inhibit prostanoid-induced smooth muscle contraction and therefore may be used in the treatment of dysmenorrhoea, premature labour, asthma and  
25 bronchitis.

The compounds of formula (I) can be used as alternative to conventional non-steroidal anti-inflammatory drugs, particularly where such non-steroidal anti-inflammatory drugs may be contraindicated such as the treatment of patients with  
30 gastrointestinal disorders including peptic ulcers, gastritis, regional enteritis, ulcerative colitis, diverticulitis, Crohn's disease, inflammatory bowel syndrome and irritable bowel syndrome, gastrointestinal bleeding and coagulation disorders, kidney disease (e.g. impaired renal function), those prior to  
35 surgery or taking anticoagulants, and those susceptible to non-steroidal anti-inflammatory drugs induced asthma.

The compounds can further be used to treat inflammation in

diseases such as vascular diseases, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anaemia, Hodgkin's disease, scleroderma, type I diabetes, myasthenia gravis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, 5 polymyositis, hypersensitivity, conjunctivitis, gingivitis, myocardial ischaemia and stroke.

Compounds of the present invention are inhibitors of cyclooxygenase-2 enzyme and are thereby useful to treat the cyclooxygenase-2 mediated diseases enumerated above. These 10 compounds can further be used for the prevention of neurodegenerative diseases such as Alzheimer's disease.

Accordingly, the 2-phenylpyran-4-one derivatives of formula (I) and pharmaceutically acceptable salts thereof, and pharmaceutical compositions comprising such compounds and/or 15 salts thereof, may be used in a method of treatment of disorders of the human body which comprises administering to a patient requiring such treatment an effective amount of a 2-phenylpyran-4-one derivative of formula (I) or a pharmaceutically acceptable salt thereof.

20 The present invention also provides pharmaceutical compositions which comprise, as an active ingredient, at least a 2-phenylpyran-4-one derivative of formula (I) or a pharmacologically acceptable salt thereof in association with a pharmaceutically acceptable excipient such as a carrier or 25 diluent. The active ingredient may comprise 0.001% to 99% by weight, preferably 0.01% to 90% by weight of the composition depending upon the nature of the formulation and whether further dilution is to be made prior to application.

Preferably the compositions are made up in a form suitable 30 for oral, topical, nasal, inhalation, rectal, percutaneous or injectable administration.

The pharmaceutically acceptable excipients which are admixed with the active compound, or salts of such compound, to form the compositions of this invention are well-known per se 35 and the actual excipients used depend inter alia on the intended method of administering the compositions.

Compositions of this invention are preferably adapted for

injectable and per os administration. In this case, the compositions for oral administration may take the form of tablets, retard tablets, sublingual tablets, capsules or liquid preparations, such as mixtures, elixirs, syrups or suspensions, all containing the compound of the invention; such preparations may be made by methods well-known in the art.

The diluents which may be used in the preparation of the compositions include those liquid and solid diluents which are compatible with the active ingredient, together with colouring or flavouring agents, if desired. Tablets or capsules may conveniently contain between 2 and 500 mg of active ingredient or the equivalent amount of a salt thereof.

The liquid composition adapted for oral use may be in the form of solutions or suspensions. The solutions may be aqueous solutions of a soluble salt or other derivative of the active compound in association with, for example, sucrose to form a syrup. The suspensions may comprise an insoluble active compound of the invention or a pharmaceutically acceptable salt thereof in association with water, together with a suspending agent or flavouring agent.

Compositions for parenteral injection may be prepared from soluble salts, which may or may not be freeze-dried and which may be dissolved in pyrogen free aqueous media or other appropriate parenteral injection fluid.

Effective doses are normally in the range of 10-600 mg of active ingredient per day. Daily dosage may be administered in one or more treatments, preferably from 1 to 4 treatments, per day.

The invention is illustrated by the following Examples, which do not limit the scope of the invention in any way.

#### EXAMPLE 1

a) To a solution of 2-(4-fluorophenyl)-1-(4-methanesulfonylphenyl)ethanone (1 g; 3.4 moles) in glacial acetic acid (15 ml), polyphosphoric acid (10 g) was added and then heated at 140°C for 16 hours. After cooling, the reaction was poured into ice-water, extracted with ethyl acetate (2 x 50



ml), the organic solution dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed under reduced pressure. The residual oil was purified by column chromatography with silica gel and ethyl acetate as eluent. 3-(4-fluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one was obtained (0.5 g) m.p.  $237^\circ\text{C}$  (Compound 2 in Table 4).

#### EXAMPLE 2

a) To a solution of 2,4-difluorophenol (3.71 g; 29 mmol) and 2-bromo-1-(4-methylsulfonylphenyl)ethanone (7.00 g; 29 mmol) in methylene chloride (50 ml) was added a solution of potassium carbonate (5.91 g; 43 mmol) and tetrabutylammonium hydrogensulfate (0.48 g; 1.4 mmol) in water (20 ml). The mixture was stirred at room temperature for 16 hours. Water (100 ml) was added, the organic phase was decanted, and the basic phase was extracted with methylene chloride (2 x 100 ml). The organic solution was dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed under reduced pressure. The resulting solid was washed with ethyl ether. 2-(2,4-Difluorophenoxy)-1-(4-methylsulfonylphenyl)ethanone was obtained (6.60 g), m.p.  $70-71^\circ\text{C}$ .

b) To a solution of 2-(2,4-difluorophenoxy)-1-(4-methylsulfonylphenyl)ethanone (6.60 g; 22 mmol) in methylene chloride (100 ml), water (20 ml) and 80% magnesium monoperoxyphosphate hexahydrate (15.26 g; 25 mmol) were added. The mixture was stirred at room temperature for 16 hours. The reaction was poured into saturated solution of sodium bicarbonate (200 ml) and extracted with methylene chloride (3 x 100 ml). The organic phase was dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed under reduced pressure. 2-(2,4-Difluorophenoxy)-1-(4-methanesulfonylphenyl)ethanone was obtained (4.97 g) as a solid, m.p.  $161-163^\circ\text{C}$ .

c) To a solution of 2-(2,4-Difluorophenoxy)-1-(4-methanesulfonylphenyl)ethanone (4.60 g; 14 mmol) in acetic acid (70 ml), polyphosphoric acid (45 g) was added and then heated at  $140^\circ\text{C}$  for 5 hours. After cooling, the reaction was poured into ice-water, extracted with ethyl acetate (2 x 100 ml), the organic solution dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed under reduced pressure. The residual oil was purified by column

chromatography with silica gel and ethyl acetate/n-hexane (1:2) as eluent. Recrystallization from ethanol gave 3-(2,4-difluorophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one (0.64 g), m.p. 191°C (Compound 45 in Table 4).

5

EXAMPLE 3

a) A solution of 4-(dibenzylsulfamoyl)benzoic acid (24 g; 63 mmol) in thionyl chloride (50 ml) was boiled under reflux for 2.5 hours and the excess of thionyl chloride removed under reduced pressure. 4-(dibenzylsulfamoyl)benzoyl chloride (25 g) was obtained as an oil which was used in the next step without purification.

b) To a solution of N,O-dimethylhydroxylamine hydrochloride (7.37 g; 75.6 mmol) and triethylamine (21.8 ml; 151 mmol) in methylene chloride (150 ml), another solution of 4-(dibenzylsulfamoyl)benzoyl chloride (25 g) in methylene chloride (150 ml) was slowly added and the resulting mixture stirred at room temperature for 16 hours. The solid was filtered off, the solvent removed under reduced pressure and the residual oil purified by column chromatography with silica gel and n-hexane-ethyl acetate 1:1 as eluent. N,O-dimethylamide of 4-(dibenzylsulfamoyl)benzoic acid (22 g) was obtained, m.p. 75°C.

c) To a suspension of magnesium (2 g; 82.4 mmol) in anhydrous tetrahydrofuran (20 ml), another solution of benzyl chloride (10.4 g; 82.4 mmol) in anhydrous tetrahydrofuran (100 ml) was slowly added. When the reaction was completed, a solution of N,O-dimethylamide of 4-(dibenzylsulfamoyl)benzoic acid (7 g; 16.5 mmol) in anhydrous tetrahydrofuran (50 ml) was slowly added while the temperature was maintained at 0°C. After stirring at the same temperature for half an hour, the reaction mixture was poured into an ammonium chloride saturated solution (100 ml), extracted with ethyl ether (3 x 75 ml) and the organic extracts dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was removed under reduced pressure and the residual oil was purified by column chromatography with silica gel and n-hexane-ethyl acetate 1:3 as eluent. N,N-dibenzyl-4-

phenylacetylbenzenesulfonamide (9.4 g) was obtained, m.p. 143°C.

d) To a solution of the above compound obtained in c) (9.4 g; 20.7 mmoles) in glacial acetic acid (140 ml), polyphosphoric acid (94 g) was added and the resulting mixture  
5 heated to 140°C for 16 hours. After cooling, the reaction mixture was poured into ice-water, extracted with ethyl acetate (3 x 150 ml) and the organic solution dried ( $\text{Na}_2\text{SO}_4$ ). The solvent was removed in vacuo and to the residual oil, concentrated sulfuric acid (38 ml) was added, then stirred at  
10 0°C for 10 minutes, further 60 minutes at room temperature and poured into ice-water. The precipitated solid was collected by filtration and purified by column chromatography with silica gel and ethyl acetate as eluent. 4-(6-methyl-4-oxo-3-phenyl-4H-pyran-2-yl)benzenesulfonamide (1.5 g) was obtained, m.p.  
15 218°C (Compound 54 in Table 4).

#### EXAMPLE 4

a) To a solution of 3,4-dichlorophenylacetophenone (5.3 g; 20 mmoles) in glacial acetic acid (90 ml), polyphosphoric  
20 acid (64 g) was added and the resulting solution heated at 140°C for 24 hours. After cooling, the reaction mixture was poured into ice-water, extracted with ethyl acetate (3 x 75 ml), the organic solution dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed under reduced pressure. The obtained residue was purified by  
25 column chromatography with silica gel and n-hexane-ethyl acetate 3:2 as eluent. 3-(3,4-dichlorophenyl)-2-phenyl-6-methylpyran-4-one (1.68 g) was obtained, m.p. 104°C.

b) A solution of the compound obtained above (1.4 g; 4.3 mmoles) in chlorosulfonic acid (12 ml) was heated at 70°C  
30 for 1.5 hours and after cooling, the reaction mixture was slowly poured into ice-water and extracted with ethyl acetate (2 x 50 ml). The organic solution was dried ( $\text{Na}_2\text{SO}_4$ ), the solvent removed under reduced pressure and to the residual oil, previously solved in methanol (10 ml), a saturated solution of  
35 ammoniac in methanol (40 ml) was slowly added. After stirring at room temperature for 1 hour, the solvent was removed under reduced pressure, the residue solved in ethyl acetate (100 ml)

and the resulting solution was washed with water (2 x 100 ml), dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent removed under reduced pressure. The residual oil was purified by column chromatography with silica gel and n-hexane-ethyl acetate 1:1 as eluent. 4-[3-(3,4-dichlorophenyl)-6-methyl-4-oxo-4H-pyran-2-yl]benzenesulfonamide (0.5 g) was obtained, m.p. 128°C (Compound 56 in Table 4).

#### EXAMPLE 5

a) To a solution of N,N-dibenzyl-4-(2-bromoacetyl)benzenesulfonamide (10.5 g, 23 mmol), and p-chlorophenol (2.94 g, 23 mmol) in methylene chloride (42 ml), potassium carbonate (4.83 g, 34.7 mmol) and tetrabutylammonium bromide (0.42 g, 1.2 mmol) in water (140 ml) was added. The reaction mixture was refluxed for 16 hours. After cooling, the mixture was diluted with methylene chloride (150 ml). The organic layer was separated, washed with water, and dried ( $\text{Na}_2\text{SO}_4$ ). The solvent was removed under reduced pressure. N,N-dibenzyl-4-[2-(4-chlorophenoxy)acetyl]benzenesulfonamide (11.7 g) was obtained as a semisolid residue, which was used in the next step without further purification.

b) To a solution of N,N-dibenzyl-4-[2-(4-chlorophenoxy)acetyl]benzenesulfonamide (11.7 g, 23 mmol) in acetic acid (105 ml), polyphosphoric acid (75 g) was added and the resulting solution was heated at 140°C for 5 hours. After cooling, the reaction mixture was poured into ice-water, extracted with ethyl acetate (3 x 150 ml), and the organic solution was dried ( $\text{Na}_2\text{SO}_4$ ). The solvent was removed under reduced pressure and the resulting oil was solved in  $\text{H}_2\text{SO}_4$  (33 ml). The mixture was stirred at room temperature for 15 minutes, and poured into ice-water. The solid was filtered off and purified by column chromatography with silica gel and ethyl acetate/methylene chloride/acetic acid (78:10:1) as eluent. 4-[3-(4-chlorophenoxy)-6-methyl-4-oxo-4H-pyran-2-yl]benzenesulfonamide (0.28 g) was obtained. m.p. 221°C (Compound 57 in Table 4).

EXAMPLE 6

a) To a solution of 3-(2,4-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one (1.7 g; 4.5 mmol) (compound 13) in dioxane (45 ml), selenium dioxide (2.2 g; 20 mmol) was added and the mixture heated in a pressure vessel at 180°C for 1 hour. After cooling, the reaction mixture was filtered, the solvent removed under reduced pressure and the residual oil purified by column chromatography with silica gel and ethyl acetate as eluent. 5-(2,4-difluorophenyl)-6-(4-methanesulfonylphenyl)-4-oxo-4H-pyran-2-carbaldehyde (0.85 g) was obtained.

b) To a solution of the above compound (0.8 g; 2.1 mmol) in formic acid (6 ml), hydroxylamine hydrochloride (0.17 g; 2.7 mmol) was added and the mixture heated at 100°C for 2 hours. After cooling, the reaction mixture was poured into ice, 2N sodium hydroxide was added until pH=7 and extracted with ethyl acetate (2 x 50 ml). The organic solution was dried (Na<sub>2</sub>SO<sub>4</sub>) the solvent removed in vacuo and the residue was dissolved in acetic anhydride (15 ml) and heated at 150°C for 3 hours. The solvent was removed under reduced pressure, the residue treated with methylene chloride (50 ml) and the resulting solution washed with 2N sodium hydroxide (2 x 25 ml). The organic solution was dried (Na<sub>2</sub>SO<sub>4</sub>), the solvent removed in vacuo and the residue purified by column chromatography with silica gel and n-hexane-ethyl acetate 1:1 as eluent. 5-(2,4-difluorophenyl)-6-(4-methanesulfonylphenyl)-4-oxo-4H-pyran-2-carbonitrile (0.2 g), m.p. 113°C (Compound 59 in Table 4).

EXAMPLE 7

a) To a solution of 3-(4-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one (4.0 g, 10.6 mmol) in dioxane (50 ml), selenium dioxide (5.9 g, 53 mmol) was added and heated into a sealed tube at 180°C for 30 minutes. After cooling, the raw material was filtered through Celite and the solvent was removed under reduced pressure. The resulting oil was purified by column chromatography with silica gel and ethyl acetate/n-hexane (2:1) as eluent. 5-(4-chlorophenyl)-6-

(4-methanesulfonylphenyl)-4-oxo-4H-pyran-2-carbaldehyde (1.80 g) was obtained.

b) To a solution of 5-(4-chlorophenyl)-6-(4-methanesulfonylphenyl)-4-oxo-4H-pyran-2-carbaldehyde (1.8 g, 4.6 mmol) in methanol (30 ml), sodium borohydride (0.26 g, 6.9 mmol) was slowly added at 0°C. The resulting mixture was stirred for 30 minutes at room temperature. The reaction mixture was concentrated and the residue was solved in ethyl acetate. The organic layer was washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was removed under reduced pressure. The resulting residue was purified by column chromatography with silica gel and ethyl acetate/n-hexane (1:1) as eluent. 3-(4-chlorophenyl)-6-hydroxymethyl-2-(4-methanesulfonylphenyl)pyran-4-one (0.9 g) was obtained. m.p. 120°C. (Compound 60 in Table 4).

#### EXAMPLE 8

a) To a solution of 3-(4-chlorophenyl)-6-hydroxymethyl-2-(4-methanesulfonylphenyl)pyran-4-one (0.5 g, 1.3 mmol) in methylene chloride (10 ml), methyl iodide (0.24 ml, 3.86 mmol), and a solution of sodium hydroxide (0.41 g, 10.3 mmol) and tetrabutylammonium chloride (50 ml) in water (0.8 ml) were added. The reaction mixture was stirred at room temperature for 18 hours. The organic layer was diluted with methylene chloride (20 ml), washed with water and dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was removed under reduced pressure. The resulting solid was purified by column chromatography with silica gel and ethyl acetate as eluent. 3-(4-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methoxymethylpyran-4-one (0.15 g) was obtained. m.p. 162°C (Compound 63 in Table 4).

#### EXAMPLE 9

a) To a solution of silver nitrate (0.88 g, 5.1 mmol) in water (4 ml), a solution of sodium hydroxide (0.42 g, 6.2 mmol) in water (4 ml) was added. The reaction mixture was stirred for 15 minutes at room temperature, and a solution 5-(4-chlorophenyl)-6-(4-methanesulfonylphenyl)-4-oxo-4H-pyran-2-


carbaldehyde in tetrahydrofuran (10 ml) was added. The reaction mixture was stirred for 3 hours at room temperature and filtered through Celite. The solvent was removed under reduced pressure and the residue was solved in ethyl acetate. The organic layer was washed with water and dried ( $\text{Na}_2\text{SO}_4$ ). The solvent was removed under reduced pressure. The resulting solid was purified by column chromatography with silica gel and ethyl acetate/methylene chloride/acetic acid (78:10:1) as eluent. 5-(4-chlorophenyl)-6-(4-methanesulfonylphenyl)-4-oxo-4H-pyran-2-carboxylic acid (0.13 g) was obtained. m.p. 236°C (Compound 65 in Table 4).

#### EXAMPLE 10

a) To a solution of 5-(4-chlorophenyl)-6-(4-methanesulfonylphenyl)-4-oxo-4H-pyran-2-carbaldehyde (0.74 g, 1.9 mmol) in methylene chloride (10 ml), diethylaminosulfide DAST (0.61 g, 3.8 mmol) was slowly added at 0°C. The reaction mixture was stirred at this temperature for 1 hour and at room temperature for 16 hours. The mixture was diluted with methylene chloride (10 ml). The organic phase was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ), and the solvent was removed under reduced pressure. The resulting residue was purified by column chromatography with silica gel and ethyl acetate/n-hexane (1:1) as eluent. 3-(4-chlorophenyl)-6-difluoromethyl-2-(4-methanesulfonylphenyl)pyran-4-one (0.1 g) was obtained. m.p. 168-170°C (Compound 67 in Table 4).

The 2-phenylpyran-4-one derivatives of general formula (I) included in Table 4 were prepared according to the processes disclosed in these Examples, but with the appropriate starting materials.

TABLE 4



Compound	R1	X	R2	R3	Method Example	Melting Point (°C)
1	CH <sub>3</sub>	single bond	C <sub>6</sub> H <sub>5</sub>	CH <sub>3</sub>	1	185
2	CH <sub>3</sub>	"	4-FC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	237
3	CH <sub>3</sub>	"	3-FC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	182
4	CH <sub>3</sub>	"	2-FC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	136-137
5	CH <sub>3</sub>	"	4-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	182
6	CH <sub>3</sub>	"	3-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	131
7	CH <sub>3</sub>	"	2-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	148
8	CH <sub>3</sub>	"	4-BrC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	198
9	CH <sub>3</sub>	"	3-BrC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	178
10	CH <sub>3</sub>	"	4-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	205
11	CH <sub>3</sub>	"	3-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	126
12	CH <sub>3</sub>	"	2-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	91-93
13	CH <sub>3</sub>	"	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	1	172
14	CH <sub>3</sub>	"	2,3-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	187
15	CH <sub>3</sub>	"	2,4-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	208
16	CH <sub>3</sub>	"	3,4-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	207
17	CH <sub>3</sub>	"	3,5-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	210
18	CH <sub>3</sub>	"	2,5-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	183
19	CH <sub>3</sub>	"	2,6-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	206



Compound	R1	X	R2	R3	Method Example	Melting Point (°C)
20	CH <sub>3</sub>	"	2,3-diClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	200
21	CH <sub>3</sub>	"	2,4-diClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	203
22	CH <sub>3</sub>	"	3,4-diClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	156
23	CH <sub>3</sub>	single bond	2,5-diClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	230
24	CH <sub>3</sub>	"	2,6-diClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	186
25	CH <sub>3</sub>	"	6-F,2-ClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	177
26	CH <sub>3</sub>	"	2-F,4-ClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	171
27	CH <sub>3</sub>	"	4-F,2-ClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	113
28	CH <sub>3</sub>	"	4-Cl,3- CH <sub>3</sub> C <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	98-99
29	CH <sub>3</sub>	"	3-Cl,4- CH <sub>3</sub> C <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	176
30	CH <sub>3</sub>	"	3-F,4- CH <sub>3</sub> OC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	137
31	CH <sub>3</sub>	"	3-Cl,4- CH <sub>3</sub> OC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	1	116
32	CH <sub>3</sub>	"	i-C <sub>3</sub> H <sub>7</sub>	CH <sub>3</sub>	1	108
33	CH <sub>3</sub>	"	C <sub>6</sub> H <sub>11</sub> (cyclohexyl)	CH <sub>3</sub>	1	98-99
34	CH <sub>3</sub>	"	2-naphthyl	CH <sub>3</sub>	1	122-123
35	CH <sub>3</sub>	"	2-indanyl	CH <sub>3</sub>	1	169
36	CH <sub>3</sub>	"	2-tetra- hydronaphthyl	CH <sub>3</sub>	1	103
37	CH <sub>3</sub>	CH <sub>2</sub>	C <sub>6</sub> H <sub>5</sub>	CH <sub>3</sub>	1	137
38	CH <sub>3</sub>	O	C <sub>6</sub> H <sub>5</sub>	CH <sub>3</sub>	2	169
39	CH <sub>3</sub>	O	4-FC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	2	189
40	CH <sub>3</sub>	O	2-FC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	2	178

Compound	R1	X	R2	R3	Method Example	Melting Point (°C)
41	CH <sub>3</sub>	O	4-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	2	196
42	CH <sub>3</sub>	O	2-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	2	198
43	CH <sub>3</sub>	O	4-BrC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	2	188
44	CH <sub>3</sub>	O	4-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	2	183
45	CH <sub>3</sub>	O	2,4-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	2	191
46	CH <sub>3</sub>	O	3,4-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	2	194
47	CH <sub>3</sub>	O	2,5-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	2	189
48	CH <sub>3</sub>	O	2,6-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	2	169
49	CH <sub>3</sub>	O	3,4-diClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	2	177
50	CH <sub>3</sub>	O	2,6-diClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	2	170
51	CH <sub>3</sub>	O	4-Cl, 3- CH <sub>3</sub> C <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	2	183
52	CH <sub>3</sub>	O	2,3,6- triClC <sub>6</sub> H <sub>2</sub>	CH <sub>3</sub>	2	216
53	CH <sub>3</sub>	O	2,4,6- triClC <sub>6</sub> H <sub>2</sub>	CH <sub>3</sub>	2	171
54	NH <sub>2</sub>	single bond	C <sub>6</sub> H <sub>5</sub>	CH <sub>3</sub>	3	218
55	NH <sub>2</sub>	"	4-FC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	3	247
56	NH <sub>2</sub>	"	3,4-diClC <sub>6</sub> H <sub>3</sub>	CH <sub>3</sub>	4	128
57	NH <sub>2</sub>	O	4-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	5	221
58	CH <sub>3</sub>	single bond	4-ClC <sub>6</sub> H <sub>4</sub>	CN	6	189
59	CH <sub>3</sub>	"	2,4-diFC <sub>6</sub> H <sub>3</sub>	CN	6	113
60	CH <sub>3</sub>	"	4-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>2</sub> OH	7	120
61	CH <sub>3</sub>	"	4-BrC <sub>6</sub> H <sub>4</sub>	CH <sub>2</sub> OH	7	128-129
62	CH <sub>3</sub>	"	2,4-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>2</sub> OH	7	173-175

Compound	R1	X	R2	R3	Method Example	Melting Point (°C)
63	CH <sub>3</sub>	"	4-ClC <sub>6</sub> H <sub>4</sub>	CH <sub>2</sub> O- CH <sub>3</sub>	8	162
64	CH <sub>3</sub>	"	2,4-diFC <sub>6</sub> H <sub>3</sub>	CH <sub>2</sub> O- CH <sub>3</sub>	8	184
65	CH <sub>3</sub>	"	4-ClC <sub>6</sub> H <sub>4</sub>	COOH	9	236
66	CH <sub>3</sub>	"	2,4-diFC <sub>6</sub> H <sub>3</sub>	COOH	9	241
67	CH <sub>3</sub>	"	4-ClC <sub>6</sub> H <sub>4</sub>	CF <sub>2</sub> H	10	168-170

Examples 11 and 12 illustrate pharmaceutical compositions according to the present invention and procedure for their preparation.

#### EXAMPLE 11

25,000 capsules each containing 100 mg of 3-(4-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one (active ingredient) were prepared according to the following formulation:

Active ingredient	2.5 Kg
Lactose monohydrate	5 Kg
Colloidal silicone dioxide	0.05 Kg
Corn starch	0.5 Kg
Magnesium stearate	0.1 Kg

Procedure

The above ingredients were sieved through a 60 mesh sieve, and were loaded into a suitable mixer and filled into 25,000 gelatine capsules.

EXAMPLE 12

100,000 Tablets each containing 50 mg of the 3-(2,4-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one (active ingredient) were prepared from the following formulation:


Active ingredient	5	Kg
Spray dried lactose	19.9	Kg
Microcrystalline cellulose	3.9	Kg
Sodium stearyl fumarate	0.2	Kg
Colloidal silicon dioxide	0.2	Kg
Carboxymethyl starch	0.8	Kg

Procedure

All the powders were passed through a screen with an aperture of 0.6 mm, then mixed in a suitable mixer for 20 minutes and compressed into 300 mg tablets using 9 mm disc and flat bevelled punches. The disintegration time of the tablets was about 3 minutes.

CLAIMS

1. A compound of formula (I):



wherein:

$R^1$  represents an alkyl or  $-NR^4R^5$  group, wherein  $R^4$  and  $R^5$  each independently represents a hydrogen atom or an alkyl group;

$R^2$  represents an alkyl,  $C_3$ - $C_7$  cycloalkyl, pyridyl, thienyl, naphthyl, tetrahydronaphthyl or indanyl group, or a phenyl group which may be unsubstituted or substituted by one or more halogen atoms or alkyl, trifluoromethyl, hydroxy, alkoxy, methylthio, amino, mono- or dialkylamino, hydroxyalkyl or hydroxycarbonyl groups;

$R^3$  represents a methyl, hydroxymethyl, alkoxymethyl,  $C_3$ - $C_7$  cycloalkoxymethyl, benzyloxymethyl, hydroxycarbonyl, nitrile, trifluoromethyl or difluoromethyl group or a  $CH_2-R^6$  group wherein  $R^6$  represents an alkyl group; and

X represents a single bond, an oxygen atom, a sulfur atom or a methylene group;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 wherein  $R^1$  represents an unsubstituted alkyl group or  $NH_2$ ,  $R^2$  represents a branched alkyl,  $C_3$ - $C_7$  cycloalkyl, naphthyl, tetrahydronaphthyl or indanyl group, an unsubstituted phenyl group or a phenyl group substituted by one or more halogen atoms, alkyl groups and/or alkoxy groups,  $R^3$  represents an unsubstituted alkyl group, a nitrile group, a hydroxymethyl group, a

methoxymethyl group, a difluoromethyl group or a hydroxycarbonyl group and X represents a single bond, an oxygen atom or a methylene group.

3. A compound according to claim 1 or 2 wherein  $R^1$  represents a methyl group,  $R^2$  represents an unsubstituted phenyl group or a phenyl group substituted by 1, 2 or 3 substituents independently selected from halogen atoms, methoxy groups and methyl groups and  $R^3$  represents a methyl group, methoxymethyl group or difluoromethyl group.

4. A compound according to any one of claims 1 to 3 wherein  $R^2$  represents a phenyl group substituted by 1, 2 or 3 substituents independently selected from halogen atoms, methoxy groups and methyl groups, one of the substituents being on the 4-position.


5. A compound according to any one of claims 1 to 3 wherein  $R^2$  represents a phenyl group substituted by one or two halogen atoms at least one of which is on the 4-position or the 2-position.

6. 3-(4-fluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(2-fluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(4-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(4-bromophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(2,4-difluorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(3,4-dichlorophenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
3-(3-chloro-4-methylphenyl)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
2-(4-methanesulfonylphenyl)-6-methyl-3-phenoxy-pyran-4-one,

3-(4-fluorophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
 3-(2-fluorophenoxy)-2-(methanesulfonylphenyl)-6-methylpyran-4-one,  
 3-(4-chlorophenoxy)-2-(methanesulfonylphenyl)-6-methylpyran-4-one,  
 3-(2-chlorophenoxy)-2-(methanesulfonylphenyl)-6-methylpyran-4-one,  
 3-(4-bromophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
 2-(4-methanesulfonylphenyl)-6-methyl-3-(4-methylphenoxy)pyran-4-one,  
 3-(2,4-difluorophenoxy)-2-(4-methanesulfonylphenyl)-6-methylpyran-4-one,  
 3-(2,5-difluorophenoxy)-2-(methanesulfonylphenyl)-6-methylpyran-4-one,  
 3-(4-chlorophenyl)-2-(4-methanesulfonylphenyl)-6-methoxymethylpyran-4-one,  
 3-(4-chlorophenyl)-6-difluoromethyl-2-(4-methanesulfonylphenyl)pyran-4-one,  
 and pharmaceutically acceptable salts thereof.

7. A process for the preparation of a compound of formula (I) as defined in any one of the preceding claims which process comprises:


(a) wherein  $R^1$  is an alkyl or  $-NR^4R^5$  group in which  $R^4$  and  $R^5$  each independently is an alkyl group,  $R^3$  is a methyl group and  $R^2$  and  $X$  are as defined in any one of claims 1 to 5, reacting a carbonyl derivative of formula (III):



wherein  $R^{1a}$  is an alkyl or a  $-NR^{4a}R^{5a}$  group in which  $R^{4a}$  and  $R^{5a}$


are each independently alkyl groups and  $R^2$  and X are as defined in any one of claims 1 to 5, with an excess of anhydrous acetic acid and polyphosphoric acid at a temperature from 100°C to 150°C;

(b) wherein  $R^1$  is an alkyl group,  $R^3$  is a methyl group, X is as defined in any one of claims 1 to 5 with the proviso that it is other than a sulfur atom and  $R^2$  is as defined in any one of claims 1 to 5, reacting a mercapto derivative of formula (VIII):




wherein  $R^{1b}$  is an alkyl group,  $X^b$  is as defined for X in any one of claims 1 to 5 with the proviso that it is other than a sulfur atom and  $R^2$  is as defined in any one of claims 1 to 5, with an oxidizing agent;

(c) wherein  $R^1$  is a  $-NR^4R^5$  group,  $R^3$  is a methyl group and  $R^2$ ,  $R^4$ ,  $R^5$  and X are as defined in any one of claims 1 to 5, reacting a chlorosulfonyl derivative of formula (X):




wherein  $R^2$  and X are as defined in any one of claims 1 to 5, with an amine of formula (XI):






wherein  $R^4$  and  $R^5$  are as defined in any one of claims 1 to 5;  
or

(d) wherein  $R^1$  is a  $-NR^4R^5$  group wherein  $R^4$  and  $R^5$  are hydrogen,  $R^3$  is a methyl group and  $R^2$  and X are as defined in any one of claims 1 to 5, by debenzoylation of the corresponding N,N-dibenzyl derivative of formula (XIV):



wherein  $R^2$  and X are as defined in any one of claims 1 to 5.

8. A compound of formula (III):




wherein  $R^{1a}$  is an alkyl or  $-NR^{4a}R^{5a}$  group in which  $R^{4a}$  and  $R^{5a}$  are each independently alkyl groups, and  $R^2$  and X are as defined in any one of claims 1 to 5.

9. A compound according to claim 8 wherein  $R^{1a}$  is a methyl group, X is an oxygen atom and  $R^2$  is selected from phenyl, 4-fluorophenyl, 2-fluorophenyl, 4-chlorophenyl, 2-chlorophenyl, 4-bromophenyl, 4-methylphenyl, 2,4-difluorophenyl, 3,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 3,4-dichlorophenyl, 2,6-dichlorophenyl, 4-chloro-3-methylphenyl, 2,3,6-trichlorophenyl and 2,4,6-

trichlorophenyl.

10. A compound of formula (VI):




wherein  $R^{1a}$  is an alkyl or  $-NR^{4a}R^{5a}$  group in which  $R^{4a}$  and  $R^{5a}$  are each independently alkyl groups,  $X^b$  is as defined for  $X$  in any one of claims 1 to 5 with the proviso that it is other than a sulfur atom and  $R^2$  is as defined in any one of claims 1 to 5.

11. A compound according to claim 10 wherein  $R^{1a}$  is a methyl group,  $X$  is an oxygen atom and  $R^2$  is selected from phenyl, 4-fluorophenyl, 2-fluorophenyl, 4-chlorophenyl, 2-chlorophenyl, 4-bromophenyl, 4-methylphenyl, 2,4-difluorophenyl, 3,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 3,4-dichlorophenyl, 2,6-dichlorophenyl, 4-chloro-3-methylphenyl, 2,3,6-trichlorophenyl and 2,4,6-trichlorophenyl.

12. Use of a compound as defined in any one of claims 8 to 11 in the preparation of a compound of formula (I) as defined in any one of claims 1 to 6.

13. Use of a compound of formula (XVI):



wherein  $R^1$  and  $R^2$  are as defined in any one of claims 1 to 5,

in the preparation of a compound of formula (I) as defined in any one of claims 1 to 5 wherein  $R^3$  is other than a methyl group.

14. A pharmaceutical composition comprising a compound according to any one of claims 1 to 6 or pharmaceutically acceptable salt thereof in admixture with a pharmaceutically acceptable carrier or diluent.

15. A compound according to any one of claims 1 to 6 or a composition according to claim 14 for use in a method of treatment of the human or animal body by therapy.

16. Use of a compound according to any one of claims 1 to 6 or a composition according to claim 14 for the manufacture of a medicament for use in the treatment of pain, fever or inflammation, to inhibit prostanoid-induced smooth muscle contraction or for the prevention of colorectal cancer or neurodegenerative diseases.

17. A method for treating pain, fever or inflammation, inhibiting prostanoid-induced smooth muscle contraction or preventing colorectal cancer or neurodegenerative diseases which comprises administering to a human or animal subject in need of treatment an effective amount of a compound according to any one of claims 1 to 6 or a composition according to claim 14.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/06873

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D309/38 C07D309/40 C07D405/04 C07D409/04 A61K31/35  
C07C317/24 C07C311/29

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 14014 A (PARKE,DAVIS &CY.) 26 May 1995 (1995-05-26) claims; example 80 ---	1,14-16
X	WO 96 06840 A (MERCK FROSST) 7 March 1996 (1996-03-07) page 37 -page 50; examples 1-11 ---	8-11
X	US 3 901 908 A (K. FITZI) 26 August 1975 (1975-08-26) column 15 -column 21; example 12 --- -/--	8-11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 January 2000

Date of mailing of the international search report

01/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/06873

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 117, no. 74, 1992 Columbus, Ohio, US; abstract no. 202023e, page 719; XP002127319 abstract & JP 04 173391 A (SANYO) 22 June 1992 (1992-06-22) ---	8-11
X	CHEMICAL ABSTRACTS, vol. 84, no. 9, 1976 Columbus, Ohio, US; abstract no. 59566b, page 538; XP002127320 abstract & CZ 156 247 A (M. PROTIVA ET AL.) 15 December 1974 (1974-12-15) ---	8-11
X	V. VALENTA ET AL.: "ALPHA-(4-TOLYL)DOPAMINE DERIVATIVES" COLLECTION OF CZECHOSLOVAK CHEMICAL COMMUNICATIONS., vol. 48, no. 5, 1983, pages 1447-1464, XP002127318 ACADEMIC PRESS, LONDON., GB ISSN: 0010-0765 page 1449 -page 1455; figure XV ---	8-11
X,P	WO 99 25697 A (KOWA CO) 27 May 1999 (1999-05-27) -& CHEMICAL ABSTRACTS, vol. 131, no. 1, 1999 Columbus, Ohio, US; abstract no. 05263, XP002127321 abstract * CAS RN 225668-35-9: ETHANONE, 2-p-CHLOROPHENYL-1-(4 -METHYLTHIO-PHENYL)-* -----	8-11

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/06873

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 17  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim 17  
is directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/06873

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9514014 A	26-05-1995	US 5808062 A	15-09-1998
		AU 695088 B	06-08-1998
		AU 4097497 A	08-01-1998
		AU 687465 B	26-02-1998
		AU 8091194 A	06-06-1995
		AU 682417 B	02-10-1997
		AU 8127694 A	06-06-1995
		BG 100564 A	31-12-1996
		CA 2174124 A	26-05-1995
		CA 2176044 A	26-05-1995
		CZ 9601343 A	14-08-1996
		EP 0729465 A	04-09-1996
		EP 0729466 A	04-09-1996
		FI 962019 A	20-05-1996
		FI 962020 A	31-05-1996
		HR 940938 A	30-06-1997
		HR 940939 A	30-06-1997
		HU 74888 A	28-02-1997
		HU 77719 A	28-07-1998
		JP 9505293 T	27-05-1997
		JP 9505295 T	27-05-1997
		MD 960171 A	30-09-1997
		NO 962015 A	15-05-1996
		NO 962016 A	15-05-1996
		NZ 275325 A	30-08-1999
		PL 314484 A	16-09-1996
		SK 63996 A	06-11-1996
		WO 9514013 A	26-05-1998
		ZA 9409147 A	21-07-1995
		ZA 9409150 A	31-07-1995
WO 9606840 A	07-03-1996	US 5521213 A	28-05-1996
		AU 689302 B	26-03-1998
		AU 3249295 A	22-03-1996
		CA 2197895 A	07-03-1996
		EP 0778834 A	18-06-1997
		JP 10504829 T	12-05-1998
US 3901908 A	26-08-1975	NONE	
JP 04173391 A	22-06-1992	NONE	
CZ 156247 A		NONE	
WO 9925697 A	27-05-1999	JP 11152274 A	08-06-1999
		AU 9762698 A	07-06-1999